

**Sobia Kauser¹, Gillian E. Westgate^{1,2},
Martin R. Green³ and
Desmond J. Tobin¹**

¹Centre for Skin Sciences, School of Life Sciences, University of Bradford, Bradford, West Yorkshire, UK; ²Westgate Consultancy Ltd, Stevington, Bedford, UK and ³Unilever R&D Colworth Laboratory, Sharnbrook, Bedford, UK
E-mail: d.tobin@bradford.ac.uk

SUPPLEMENTARY MATERIAL

Supplementary material is linked to the online version of the paper at <http://www.nature.com/jid>

REFERENCES

- Arck PC, Overall R, Spatz K *et al.* (2006) Towards a "free radical theory of graying": melanocyte apoptosis in the aging human hair follicle is an indicator of oxidative stress induced tissue damage. *FASEB J* 20:1567–9
- Commo S, Gaillard O, Thibaut S *et al.* (2004) Absence of TRP-2 in melanogenic melanocytes of human hair. *Pigment Cell Res* 17:488–97
- Gunn DA, Rexbye H, Griffiths CE *et al.* (2009) Why some women look young for their age. *PLoS One* 4:e8021
- Michard Q, Commo S, Rocchetti J *et al.* (2008) TRP-2 expression protects HEK cells from dopamine- and hydroquinone-induced toxicity. *Free Radic Biol Med* 45:1002–10
- Nishimura EK, Granter SR, Fisher DE (2005) Mechanisms of hair graying: incomplete melanocyte stem cell maintenance in the niche. *Science* 307:720–4
- Pawelek JM, Lerner AB (1978) 5,6-Dihydroxyindole is a melanin precursor showing potent cytotoxicity. *Nature* 276:626–8
- Pelle E, Mammone T, Maes D *et al.* (2005) Keratinocytes act as a source of reactive oxygen species by transferring hydrogen peroxide to melanocytes. *J Invest Dermatol* 124:793–7
- Reynolds A, Murray PI, Colloby PS (1998) Darkening of eyelashes in a patient treated with latanoprost. *Eye (Lond)* 12(Part 4):741–3
- Shaffrali FC, McDonagh AJ, Messenger AG (2002) Hair darkening in porphyria cutanea tarda. *Br J Dermatol* 146:325–9
- Slominski A, Tobin DJ, Shibahara S *et al.* (2004) Melanin pigmentation in mammalian skin and its hormonal regulation. *Physiol Rev* 84:1155–228
- Thibaut S, De Becker E, Caisey L *et al.* (2009) Human eyelash characterization. *Br J Dermatol* 162:304–10
- Tobin DJ, Paus R (2001) Graying: gerontobiology of the hair follicle pigmentary unit. *Exp Gerontol* 36:29–54
- Van Raamsdonk CD, Barsh GS, Wakamatsu K *et al.* (2009) Independent regulation of hair and skin color by two G protein-coupled pathways. *Pigment Cell Melanoma Res* 22:819–26
- Wood JM, Decker H, Hartmann H *et al.* (2009) Senile hair graying: H₂O₂-mediated oxidative stress affects human hair color by blunting methionine sulfoxide repair. *FASEB J* 23:2065–75
- Wood JM, Gibbons NC, Chavan B *et al.* (2008) Computer simulation of heterogeneous single nucleotide polymorphisms in the catalase gene indicates structural changes in the enzyme active site, NADPH-binding and tetramerization domains: a genetic predisposition for an altered catalase in patients with vitiligo? *Exp Dermatol* 17:366–71

Full Sequencing of the *FLG* Gene in Italian Patients with Atopic Eczema: Evidence of New Mutations, but Lack of an Association

Journal of Investigative Dermatology (2011) **131**, 982–984; doi:10.1038/jid.2010.398; published online 3 February 2011

TO THE EDITOR

Atopic eczema (AE) (OMIM %603165) is the most common chronic inflammatory skin disease, characterized by xerosis, pruritus, and erythematous lesions with increased transepidermal water loss. In recent years, it has been suggested that the epidermal skin barrier has a significant role in AE disease susceptibility and severity (Smith *et al.*, 2006; Cork *et al.*, 2009). Sandilands *et al.* (2006) demonstrated that null mutations within the filaggrin gene (*FLG*) strongly predispose individuals to AE. Two *FLG* null alleles (R501X and 2282del4) have been shown to be significantly associated with AE in several European populations (Palmer *et al.*, 2006; Weidinger *et al.*, 2007). Recently, a meta-analysis of the most common *FLG*

variants in European populations, involving 5,791 eczema cases and 26,454 controls (Rodriguez *et al.*, 2009), revealed that there is a high risk conferred by R501X and 2282del4 across the studies, with an overall odds ratio of 3.14 and 2.78, respectively. Indeed, large differences in carrier frequencies exist across Europe, ranging from 1.4% in an Italian population (Giardina *et al.*, 2008) to 63% in an Irish population (Palmer *et al.*, 2006). Recently, we observed that in Italian patients the frequencies of R501X and 2282del4 are strongly reduced with respect to those described in other patients of European origin, and the frequencies are similar between cases and controls (0.6 vs. 0.0% and 0.9 vs. 0.5%, respectively). In order to determine whether other

mutations located elsewhere in *FLG* confer risk to AE, we performed a full sequencing of *FLG* in Italian patients.

We performed a sequencing of the full *FLG* gene in a cohort of 220 Italian AE patients (recruited by IDI-Istituto Dermatologico dell'Immacolata and Fatebenefratelli Hospital). We then determined the frequency of variations and mutations in a cohort of 201 healthy subjects. The diagnosis of AE in our case cohort was made by experienced dermatologists or by a pediatric allergologist. The cohort consisted of 85% of cases with the intrinsic subtype and 15% with the extrinsic form of AE. These subtypes and severity of AE have been established based on the total IgE level (extrinsic subtype >500 ng l⁻¹) and using the scoring atopic dermatitis. Further clinical details of Italian patients are available in

Abbreviations: AE, atopic eczema; *FLG*, filaggrin gene; LD, linkage disequilibrium




Figure 1. *FLG* null mutations and linkage disequilibrium (LD) plot. (a) Electropherograms of the null mutations. **(b)** Gene location of filaggrin (*FLG*) null mutations. **(c)** LD pattern in atopic eczema cases. **(d)** LD pattern in healthy controls.

Bergboer *et al.* (2010). Written consent was obtained from all of the patients or their parents. The study was approved by the ethics committee of each participating institute and complied with the guidelines of the Declaration of Helsinki.

Genomic DNA was extracted from peripheral blood using standard procedures. Sequence analysis of the third exon of *FLG* using 11 overlapping fragments amplified by 11 sequence-specific primer pairs was performed

(see Supplementary Table S1 online). Furthermore, the promoter region and exon 2 were also analyzed. PCR fragments were run on an ABI 3130xl automated sequencer (Applied Biosystems, Foster City, CA). We identified

Table 1. Allele frequencies of three unreported mutations

Allele	R1798X_C>T		E3603X_G>T		R3638X_A>T	
	C	T	G	T	A	T
Cases	0.998	0.002	0.995	0.005	0.97	0.03
Controls	0.997	0.003	1	0	0.96	0.04

three null mutations that, to our knowledge, are previously unreported (R1798X (5392 C>T), E3603X (10810 G>T), and R3638X (10912 A>T)) (Figure 1a and b) and are located in the repeats 5 (R1798X) and 10 (E3603X and R3638X), adjacent to the other null mutations identified thus far. Subsequently, we assessed the frequency of the mutations in 201 healthy controls. We found that the frequency of the mutations was similar in the cases and controls (Table 1). The mutations R1798X and E3603X are rare in an Italian population, and, although not statistically significant, further studies in Italians, as well as other populations, will be needed to confirm their frequency and/or rule out a functional role. The null mutation, R3638X, has a higher frequency in healthy controls (4%) and might be considered a polymorphic variant. Moreover, we identified a number of variations, but failed to observe evidence of an association in our patients. Therefore, it is likely that in Italian cases there are other genes associated with the 1q21 region (ATOD2). In this regard, we believe that a study involving Italians, as well as other 1q21-linked samples, can be particularly useful to detect population-specific associated alleles. As in a critical region that harbors the susceptibility variant the extent of linkage disequilibrium (LD) may be different between cases and controls, we decided to contrast LD patterns between AE cases and controls (Figure 1c and d). We considered all the variations with a frequency >0.1 at the *FLG* locus and found a different pattern of LD. In particular, we observed a more conserved block between markers rs12405241 and 1432 in controls with respect to the cases. Although these data need to be replicated in other populations and using a higher

number of markers, it is suggested that disease-bearing chromosomes and control chromosomes arise from different pools, with disease-bearing chromosomes showing greater heterogeneity. Further studies will be needed to verify the existence of multiple susceptibility alleles spanning the ATOD2 locus.

Taken together, these results demonstrate that only a small percentage of Italian AE patients carry mutations in the *FLG* gene, which therefore should not be considered as a major risk factor for AE in the Italian population. The low frequency of *FLG* null alleles and the absence of association in the AE cohort may reflect the strong purifying selection of *FLG* mutations in chromosome 1 of Italian patients. In this regard, it is notable that the frequency of mutations is strongly reduced in healthy controls with respect to that observed in different European populations, further supporting the existence of a different ethnic origin of Italian population and ruling out a possible confounding effect of clinical and genetic heterogeneity characterization of samples. An alternative explanation would be that the mutations are strongly advantageous in northern Europe. As a result, the geographic distribution of the frequencies of *FLG* mutations is heterogeneous among European populations. Currently there is no evidence of *FLG* mutations in Mediterranean populations, making possible the existence of a population-specific gradient across Europe. With this perspective in mind, a pan-European study of *FLG* mutations should be conducted to further address this hypothesis.

CONFLICT OF INTEREST

The authors state no conflict of interest.

Raffaella Cascella¹, Valeria Foti Cuzzola², Tiziana Lepre¹, Elena Galli³, Viviana Moschese⁴, Loredana Chini⁴,

Cinzia Mazzanti⁵, Paola Fortugno⁶, Giuseppe Novelli^{1,7,8,9} and Emiliano Giardina¹

¹Centre of Excellence for Genomic Risk Assessment in Multifactorial and Complex Diseases, School of Medicine, Tor Vergata University of Rome, Rome, Italy; ²IRCCS Centro Neurolesi "Bonino-Pulejo", Messina, Italy; ³San Pietro Hospital, Fatebenefratelli, Rome, Italy; ⁴Department of Pediatrics, Tor Vergata University, Rome, Italy; ⁵Day Hospital Service, II Dermatology Division, Istituto Dermatologico dell'Immacolata, IDI-IRCCS, Rome, Italy; ⁶Laboratory of Molecular and Cell Biology, Istituto Dermatologico dell'Immacolata, IDI-IRCCS; ⁷Department of Cardiovascular Medicine, University of Arkansas for Medical Sciences, Little Rock, Arkansas, USA; ⁸Department of Dermatology, Tor Vergata University of Rome, Rome, Italy and ⁹Department of Applied Clinical and Medical Therapy, Rheumatological Unit, "La Sapienza" University of Rome, Rome, Italy
E-mail: emiliano.giardina@uniroma2.it

SUPPLEMENTARY MATERIAL

Supplementary material is linked to the online version of the paper at <http://www.nature.com/jid>

REFERENCES

- Bergboer JG, Zeeuwen PL, Irvine AD *et al.* (2010) Deletion of late cornified envelope 3B and 3C genes is not associated with atopic dermatitis. *J Invest Dermatol* 130:2057-61
- Cork MJ, Danby SG, Vasilopoulos Y *et al.* (2009) Epidermal barrier dysfunction in atopic dermatitis. *J Invest Dermatol* 128:1892-908
- Giardina E, Paolillo N, Sinibaldi C *et al.* (2008) RX501 and 2282del4 filaggrin mutations do not confer susceptibility to psoriasis and atopic dermatitis in Italian patients. *Dermatology* 216:83-4
- Palmer CN, Irvine AD, Terron-Kwiatkowski A *et al.* (2006) Common loss-of-function variants of the epidermal barrier protein filaggrin are a major predisposing factor for atopic dermatitis. *Nat Genet* 38:441-6
- Rodriguez E, Baurecht H, Herberich E *et al.* (2009) Meta-analysis of filaggrin polymorphisms in eczema and asthma: robust risk factors in atopic disease. *J Allergy Clin Immunol* 123:1361-70
- Sandilands A, O'Regan GM, Liao H *et al.* (2006) Prevalent and rare mutations in the gene encoding filaggrin cause ichthyosis vulgaris and predispose individuals to atopic dermatitis. *J Invest Dermatol* 126:1770-5
- Smith FJ, Irvine AD, Terron-Kwiatkowski A *et al.* (2006) Loss-of-function mutations in the gene encoding filaggrin cause ichthyosis vulgaris. *Nat Genet* 38:337-42
- Weidinger S, Rodríguez E, Stahl C *et al.* (2007) Filaggrin mutations strongly predispose to early-onset and extrinsic atopic dermatitis. *J Invest Dermatol* 127:724-6